

## DRUG FACTS

**Active ingredient(s)**

**Purpose**

### Use(s)

#### Uses

Temporarily relieves minor aches and pains associated with

- headache
- backache
- common cold
- minor arthritis pain
- toothache
- menstrual cramps
- muscular aches

Temporarily reduces fever.

### Warnings

#### Warnings

##### Do not use

##### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

##### Ask a doctor before use if you have

##### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

##### Ask a doctor or pharmacist before use if you are

##### Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

##### When using this product

##### When using this product

- take with food or milk if stomach upset occurs

##### Stop use and ask a doctor if

##### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

##### If pregnant or breast-feeding

##### Keep out of reach of children

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

#### Directions

- do not use more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

### Other information

#### Drug Facts

**Allergy alert:** Ibuprofen may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- rash
- blisters
- skin reddening
- facial swelling
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

##### Adults and children: (12 years and older)

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

##### Children under 12 years:

Do not give to children under 12 years of age.

##### Other information

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets

- do not use any opened or torn packets

***Inactive ingredients***

***Inactive ingredients***

carnauba wax<sup>\*</sup>, cellulose<sup>\*</sup>, colloidal silicon dioxide, corn starch<sup>\*</sup>, hypromellose, iron oxide red<sup>\*</sup>, lactose, magnesium stearate, microcrystalline cellulose<sup>\*</sup>, polydextrose, polyethylene glycol, povidone, silica<sup>\*</sup>, sodium lauryl sulfate<sup>\*</sup>, sodium starch glycolate, stearic acid<sup>\*</sup>, titanium dioxide, triacetin<sup>\*</sup>  
<sup>\*</sup> may contain